PRODUCT INFORMATION



1,2,3-Trilauroyl Glycerol

Item No. 23336

CAS Registry No.:	538-24-9	
Formal Name:	dodecanoic acid, 1,1',1"-(1,2,3-propanetriyl)	0
	ester	$\square \square $
Synonyms:	Glycerol Trilaurate, NSC 4061,	
	TG(12:0/12:0/12:0), Tridodecanoate,	
	Tridodecanoin, Tridodecanoyl Glycerol,	
	Trilaurin, Trilauroylglycerol	
MF:	C ₃₉ H ₇₄ O ₆	
FW:	639.0	
Purity:	≥95%	$\qquad \qquad $
Supplied as:	A crystalline solid	Ö
Storage:	-20°C	
Stability:	≥2 years	
Information represent	s the product specifications. Batch specific analytical re	esults are provided on each certificate of analysis

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

1,2,3-Trilauroyl glycerol is supplied as a crystalline solid. A stock solution may be made by dissolving the 1,2,3-trilauroyl glycerol in the solvent of choice. 1,2,3-Trilauroyl glycerol is soluble in organic solvents such as ethanol and dimethyl formamide, which should be purged with an inert gas. The solubility of 1,2,3-trilauroyl glycerol in these solvents is approximately 1 and 20 mg/ml, respectively.

Description

1,2,3-Trilauroyl glycerol is a triacylglycerol found in dietary fats.¹ 1,2,3-Trilauroyl glycerol increases HMG-CoA reductase activity and cholesterol synthesis in the jejunum of rats fed a diet containing it at 10% (w/w).² Formulations containing 1,2,3-trilauroyl glycerol are used in cosmetic products to condition skin and as thickening agents.³ 1,2,3-Trilauroyl glycerol is also used to form the lipid matrices of solid lipid nanoparticles.4

References

- 1. Litchfield, C., Miller, E., Harlow, R.D., et al. The triglyceride composition of 17 seed fats rich in octanoic, decanoic, or lauric acid. Lipids 2(4), 345-350 (1967).
- 2. Oku, H., and Sugano, M. Dietary fat dependence of intestinal 3-hydroxy-3-methylglutaryl coenzyme A (HMG-CoA) reductase activity in rats. J. Nutr. 115(7), 880-889 (1985).
- 3. Johnson, W., Jr., and C.I.R.E. Panel. Final report on the safety assessment of trilaurin, triarachidin, tribehenin, tricaprin, tricaprylin, trierucin, triheptanoin, triheptylundecanoin, triisononanoin, triisopalmitin, triisostearin, trilinolein, trimyristin, trioctanoin, triolein, tripalmitin, tripalmitolein, triricinolein, tristearin, triundecanoin, glyceryl triacetyl hydroxystearate, glyceryl triacetyl ricinoleate, and glyceryl stearate diacetate. Int. J. Toxicol. 20 (Suppl 4), 61-94 (2001).
- 4. Mohtar, N., Khan, N.A.K., and Darwis, Y. Solid lipid nanoparticles of atovaquone based on 2(4) full-factorial design. Iran J. Pharm. Res. 14(4), 989-1000 (2015).

WARNING THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFFTY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

uyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 06/04/2020

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897 [734] 971-3335 FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.CAYMANCHEM.COM